



# The scale factor: a new degree of freedom in phase-type approximation

A. Bobbio<sup>a,\*</sup>, A. Horváth<sup>b</sup>, M. Telek<sup>c</sup>

<sup>a</sup> *Dipartimento di Scienze e Tecnologie Avanzate, Università del Piemonte Orientale, Alessandria, Italy*

<sup>b</sup> *Dipartimento di Informatica, Università di Torino, Turin, Italy*

<sup>c</sup> *Department of Telecommunications, Technical University of Budapest, Budapest, Hungary*

---

## Abstract

This paper introduces a unified approach to phase-type approximation in which the discrete and the continuous phase-type models form a common model set. The models of this common set are assigned with a non-negative real parameter, the *scale factor*. The case when the scale factor is strictly positive results in discrete phase-type distributions and the scale factor represents the time elapsed in one step. If the scale factor is 0, the resulting class is the class of CPH distributions. Applying the above view, it is shown that there is no qualitative difference between the discrete and the CPH models. Based on this unified view of phase-type models one can choose the best phase-type approximation of a stochastic model by optimizing the scale factor.

© 2003 Elsevier B.V. All rights reserved.

*Keywords:* Discrete and continuous phase-type distributions; Phase-type expansion; Approximate analysis

---

## 1. Introduction

This paper presents new comparative results on the use of discrete phase-type (DPH) distributions [22] and of continuous phase-type (CPH) distributions [23] in applied stochastic modeling.

DPH distributions of order  $n$  are defined as the time to absorption in a discrete-state discrete-time Markov chain (DTMC) with  $n$  transient states and one absorbing state. CPH distributions of order  $n$  are defined, similarly, as the distribution of the time to absorption in a discrete-state continuous-time Markov chain (CTMC) with  $n$  transient states and one absorbing state. The above definition implies that the properties of a DPH distribution are computed over the set of the natural numbers while the properties of a CPH distribution are defined as a function of a continuous time variable  $t$ . When DPH distributions are used to model timed activities, the set of the natural numbers must be related to a time measure. Hence, a new parameter need to be introduced that represents the time span associated to each step. This new parameter is the scale factor of the DPH distribution, and can be viewed as a new degree of freedom,

---

\* Corresponding author.

*E-mail addresses:* andrea.bobbio@mfn.unipmn.it (A. Bobbio), horvath@di.unito.it (A. Horváth), telek@hit.bme.hu (M. Telek).

since its choice largely impacts the shape and properties of a DPH distribution over the continuous time axes. When DPH distributions are used to approximate a given continuous distribution, the scale factor affects the goodness of the fit.

The paper starts discussing to what extent DPH or CPH distributions can be utilized to fit a given continuous distribution. It is shown that a DPH distribution of any order converges to a CPH distribution of the same order as the scale factor goes to 0. Even so, the DPH class contains distributions whose behavior differs substantially from the one of the corresponding distributions in the CPH class. Two main peculiar points differentiate the DPH class from the CPH class. The first point concerns the coefficient of variation: indeed, while in the continuous case the minimum coefficient of variation is a function of the order only and its lower bound is given by the well-known theorem of Aldous and Shepp [1], in the discrete case the minimum coefficient of variation is proved to depend both on the order and on the mean (and hence on the scale factor) [25]. Furthermore, it is easy to see that for any order, there exist members of the DPH class that represent a deterministic value with a coefficient of variation equal to 0. Hence, for any order (greater than 1), the coefficient of variation of the DPH class spans from 0 to  $\infty$ .

The second peculiar point that differentiates the DPH class is the support of the distributions. While a CPH distribution (of any order) has always an infinite support, there exist members of the DPH class of any order that have a finite support (between a minimum non-negative value and a maximum) or have a mass equal to one concentrated in a single value (deterministic distribution).

It turns out that the possibility of

- tuning the scale factor to optimize the goodness of the fit,
- having distributions with coefficient of variation spanning from 0 to  $\infty$ ,
- representing deterministic values exactly,
- coping with finite support distributions,

makes the DPH class a very interesting and challenging class of distributions to be explored in applied stochastic models. The purpose of this paper is to show how these favorable properties can be exploited in practice, and to provide guidelines to the modeler to a reasonably good choice of the distributions to be used. Indeed, since a DPH distribution tends to a CPH distribution as the scale factor approaches 0, considering the scale factor as a new decision variable in a fitting experiment, and finding the value of the optimal scale factor (with respect to some error measure) provides a valuable tool to decide whether to use a discrete or a continuous approximation to the given problem.

The fitting problem for the CPH class has been extensively studied and reported in the literature by resorting to a variety of structures and numerical techniques [2,3,9,18,24]. Conversely, the fitting problem for the DPH class has received very little attention [6].

In recent years, a considerable effort has been devoted to define models with generally distributed timings [7,10] and to merge in the same model random variables and deterministic duration [21]. Analytical solutions are possible in special cases, and the approximation of the original problems by means of CPH distributions is a rather well-known technique [8,12]. This paper is aimed at emphasizing that DPH approximation may provide a more convenient alternative with respect to CPH approximation, and also to provide a way to quantitatively support this choice. Furthermore, the use of DPH approximation can be extended from stochastic models to functional analysis where time intervals with non-deterministic choice are considered [4,5]. Finally, discretization techniques for continuous problems [13,17] can be restated in terms of DPH approximations.

The rest of the paper is organized as follows. After defining the notation to be used in the paper in Section 2, Section 3 discusses the peculiar properties of the DPH class with respect to the CPH class. Some guidelines for bounding the parameters of interest and extensive numerical experiments to show how the goodness of the fit is influenced by the optimal choice of the scale factor are reported in Section 4. Section 5 discusses the quality of the approximation when passing from the analysis of a single distribution to the analysis of performance measures in complete non-Markovian stochastic models. The paper is concluded in Section 6.

## 2. Definition and notation

A DPH distribution [22,23] is the distribution of the time to absorption in a DTMC with  $n$  transient states, and one absorbing state numbered  $(n + 1)$ . The one-step transition probability matrix of the corresponding DTMC can be partitioned as

$$\hat{\mathbf{B}} = \begin{bmatrix} \mathbf{B} & \mathbf{b} \\ \mathbf{0} & 1 \end{bmatrix}, \quad (1)$$

where  $\mathbf{B} = [b_{ij}]$  is the  $(n \times n)$  matrix collecting the transition probabilities among the transient states,  $\mathbf{b} = [b_{i,n+1}]^T$  is the column vector of length  $n$  grouping the probabilities from any state to the absorbing one, and  $\mathbf{0} = [0]$  is the zero vector. The initial probability vector  $\hat{\alpha} = [\alpha, \alpha_{n+1}]$  is of length  $(n + 1)$ , with  $\sum_{j=1}^n \alpha_j = 1 - \alpha_{n+1}$ . In the present paper, we consider only the class of DPH distributions for which  $\alpha_{n+1} = 0$ , but the extension to the case when  $\alpha_{n+1} > 0$  is straightforward. The tuple  $(\alpha, \mathbf{B})$  is called the representation of the DPH distribution, and  $n$  the order.

Similarly, a CPH distribution [23] is the distribution of the time to absorption in a CTMC with  $n$  transient states, and one absorbing state numbered  $(n + 1)$ . The infinitesimal generator  $\hat{\mathbf{Q}}$  of the CTMC can be partitioned in the following way:

$$\hat{\mathbf{Q}} = \begin{bmatrix} \mathbf{Q} & \mathbf{q} \\ \mathbf{0} & 1 \end{bmatrix}, \quad (2)$$

where  $\mathbf{Q}$  is a  $(n \times n)$  matrix that describes the transient behavior of the CTMC and  $\mathbf{q}$  is the column vector grouping the transition rates to the absorbing state. Let  $\hat{\alpha} = [\alpha, \alpha_{n+1}]$  be the  $(n + 1)$  initial probability (row) vector with  $\sum_{i=1}^n \alpha_i = 1 - \alpha_{n+1}$ . The tuple  $(\alpha, \mathbf{Q})$  is called the representation of the CPH distribution, and  $n$  the order.

It has been shown in [11] for the continuous case and in [6] for the discrete case that the representations in (1) and (2), do not provide a convenient form for running a fitting algorithm because of the too high number of free parameters. Instead, resorting to acyclic phase-type distributions, the number of free parameters is reduced significantly since both in the continuous and the discrete case a canonical form can be used. Fig. 1 gives the canonical form and associated constraints for the continuous case [11]. The canonical form and its constraints for the discrete case [6] is depicted in Fig. 2. In both cases the canonical form corresponds to a “parameter-ordered” mixture of Hypo-exponential distributions.

Fitting algorithm that provides acyclic CPH distributions has been provided in [2,3]. For acyclic DPH distributions, an algorithm has been presented in [6]. Experiments suggests (an exhaustive comparison of fitting algorithms can be found in [20]) that, from the point of view of applications, it is reasonable to restrict our attention to the class of acyclic phase-type distributions.




Fig. 1. Canonical representation of acyclic CPH distributions and its constraints.




Fig. 2. Canonical representation of acyclic DPH distributions and its constraints.

### 3. Comparing properties of CPH and DPH distributions

CTMC are defined as a function of a continuous time variable  $t$ , while DTMC are defined over the set of the natural numbers. In order to relate the number of jumps in a DTMC with a time measure, a time span must be assigned to each step. Let  $\delta$  be (in some arbitrary units) the scale factor, i.e. the time span assigned to each step. The value of  $\delta$  establishes an equivalence between the sentence “probability at the  $k$ th step” and “probability at time  $k\delta$ ”, and hence, defines the time scale on which the properties of the DTMC are measured. The consideration of the scale factor  $\delta$  introduces a new parameter, and consequently a new degree of freedom, in the DPH class with respect to the CPH class. In the following, we discuss how this new degree of freedom impacts the properties of the DPH class and how it can be exploited in practice.

Let  $u$  be an “unscaled” DPH distributed random variable (r.v.) of order  $n$  with representation  $(\alpha, \mathbf{B})$ , defined over the set of the non-negative natural numbers. Let us consider a scale factor  $\delta$ ; the scaled r.v.  $\tau = \delta u$  is defined over the discrete set of time points  $(1\delta, 2\delta, 3\delta, \dots, k\delta, \dots)$ , being  $k$  a non-negative natural number. For the unscaled and the scaled DPH r.v. the following equations hold:

$$F_u(k) = \Pr\{u \leq k\} = 1 - \alpha \mathbf{B}^k \mathbf{e}, \quad F_\tau(\delta k) = \Pr\{\tau \leq \delta k\} = 1 - \alpha \mathbf{B}^k \mathbf{e},$$

$$m_u^i = E(u^i), \quad m_\tau^i = E(\tau^i) = \delta^i E(u^i), \quad i \geq 1, \tag{3}$$

where  $\mathbf{e}$  is the column vector of ones, and  $E(u^i)$  is the  $i$ th moment calculated from the factorial moments of  $u$ :  $E(u(u-1)\dots(u-i+1)) = i! \alpha (\mathbf{I} - \mathbf{B})^{-i} \mathbf{B}^{i-1} \mathbf{e}$ . It is evident from (3) that the mean  $m_\tau$  of the scaled r.v.  $\tau$  is  $\delta$  times the mean  $m_u$  of the unscaled r.v.  $u$ . While  $m_u$  is an invariant of the representation  $(\alpha, \mathbf{B})$ ,  $\delta$  is a free parameter; adjusting  $\delta$ , the scaled r.v. can assume any mean value  $m_\tau \geq 0$ . On the other hand, one can easily infer from (3) that the coefficients of variation of  $\tau$  and  $u$  are equal. A consequence of the above properties is that one can easily provide a scaled DPH of order  $\geq 2$  with arbitrary mean and arbitrary coefficient of variation with an appropriate scale factor. Or more formally: the unscaled DPH r.v.  $u$  of any order  $n > 1$  can exhibit a coefficient of variation between  $0 \leq cv_u^2 \leq \infty$ . For  $n = 1$  the coefficient of variation ranges between  $0 \leq cv_u^2 \leq 1$ .

As mentioned earlier, an important property of the DPH class with respect to the CPH class is the possibility of exactly representing a deterministic delay. A deterministic distribution with value  $a$  can be realized by means of a scaled DPH distribution with  $n$  phases with scale factor  $\delta$  if  $n = a/\delta$  is integer. In this case, the structure of the DPH distribution is such that phase  $i$  is connected with probability 1 only to phase  $i + 1$  ( $i = 1, \dots, n$ ), and with an initial probability concentrated in state 1. If  $n = a/\delta$  is not integer for the given  $\delta$ , the deterministic behavior can only be approximated.

### 3.1. First-order discrete approximation of CTMCs

Given a CTMC with infinitesimal generator  $\tilde{\mathbf{Q}}$ , the transition probability matrix over an interval of length  $\delta$  can be written as

$$e^{\tilde{\mathbf{Q}}\delta} = \sum_{i=0}^{\infty} \frac{(\tilde{\mathbf{Q}}\delta)^i}{i!} = \mathbf{I} + \tilde{\mathbf{Q}}\delta + \sigma(\delta),$$

hence the first-order approximation of  $e^{\tilde{\mathbf{Q}}\delta}$  is matrix  $\mathbf{\Pi}(\delta) = \mathbf{I} + \tilde{\mathbf{Q}}\delta$ .  $\mathbf{\Pi}(\delta)$  is a proper stochastic matrix if  $\delta < 1/q$ , where  $q = \max_{i,j} |\tilde{\mathbf{Q}}_{ij}|$ .  $\mathbf{\Pi}(\delta)$  is the exact transition probability matrix of the CTMC assumed that at most one transition occurs in the interval of length  $\delta$ .

We can approximate the behavior of the CTMC at time  $(\delta, 2\delta, 3\delta, \dots, k\delta, \dots)$  using the DTMC with transition probability matrix  $\mathbf{\Pi}(\delta)$ . The approximate transition probability matrix at time  $t = k\delta$  is

$$\mathbf{\Pi}(\delta)^k = (\mathbf{I} + \tilde{\mathbf{Q}}\delta)^{t/\delta}$$

The following theorem proves the property that the above first-order approximation becomes exact as  $\delta \rightarrow 0$ .

**Theorem 1.** *As the length of the interval of the first-order approximation,  $\delta$  tends to 0, such that  $t = k\delta$  the approximate transition probability matrix tends to the exact one.*

**Proof.** The scalar version of the applied limiting behavior is well-known in the form

$$\lim_{x \rightarrow 0} (1 + ax)^{1/x} = e^a.$$

Since matrices  $\mathbf{I}$  and  $\tilde{\mathbf{Q}}$  commute we can obtain the matrix version of the same expression as follows:

$$\begin{aligned} \lim_{\delta \rightarrow 0} (\mathbf{I} + \tilde{\mathbf{Q}}\delta)^{t/\delta} &= \lim_{k \rightarrow \infty} \mathbf{\Pi} \left( \frac{t}{k} \right)^k = \lim_{k \rightarrow \infty} \left( \mathbf{I} + \frac{\tilde{\mathbf{Q}}t}{k} \right)^k = \lim_{k \rightarrow \infty} \sum_{j=0}^k \binom{k}{j} \left( \frac{\tilde{\mathbf{Q}}t}{k} \right)^j \\ &= \lim_{k \rightarrow \infty} \sum_{j=0}^k \frac{(\tilde{\mathbf{Q}}t)^j}{j!} \frac{k!}{k^j (k-j)!} = \sum_{j=0}^{\infty} \frac{(\tilde{\mathbf{Q}}t)^j}{j!} = e^{\tilde{\mathbf{Q}}t}. \quad \square \end{aligned}$$

An obvious consequence of Theorem 1 for PH distributions is given in the following corollary.

**Corollary 1.** *Given a scaled DPH distribution of order  $n$ , representation  $(\alpha, \mathbf{I} + \mathbf{Q}\delta)$  and scale factor  $\delta$ , the limiting behavior as  $\delta \rightarrow 0$  is the CPH distribution of order  $n$  with representation  $(\alpha, \mathbf{Q})$ .*




Fig. 3.  $m_u \leq n$ .

3.2. The minimum coefficient of variation

It is known that one of the main limitation in approximating a given distribution by a PH distribution is the attainable minimal coefficient of variation,  $cv_{\min}^2$ . In order to discuss this point, we recall two theorems that state the  $cv_{\min}^2$  for the class of CPH and DPH distributions.

**Theorem 2** (Aldous and Shepp [1]). *The  $cv_{\min}^2$  of a CPH distributed r.v. of order  $n$  is  $cv_{\min}^2 = 1/n$  and is attained by the Erlang( $n$ ) distribution independent of its mean  $m_c$  or of its parameter  $\lambda = n/m_c$ .*

The corresponding theorem for the unscaled DPH class has been proved in [25]. In the following,  $\lfloor x \rfloor$  denotes the integer part and  $\langle x \rangle$  denotes the fractional part of  $x$ .

**Theorem 3.** *The  $cv_{\min}^2$  of an unscaled DPH r.v. of order  $n$  and mean  $m_u$  is*

$$\frac{\langle m_u \rangle (1 - \langle m_u \rangle)}{m_u^2} \quad \text{if } m_u \leq n, \quad \frac{1}{n} - \frac{1}{m_u} \quad \text{if } m_u > n. \tag{4}$$

The unscaled DPH r.v. which exhibits this minimal coefficient of variation has the following canonical structure:

- if  $m_u \leq n$ : each state is connected to the next with probability 1 and the non-zero initial probabilities are  $\alpha_{n-\lfloor m_u \rfloor} = \langle m_u \rangle$  and  $\alpha_{n-\lfloor m_u \rfloor+1} = 1 - \langle m_u \rangle$  (Fig. 3),
- if  $m_u > n$ : each state is connected to the next with probability  $n/m_u$  and the only non-zero initial probability is  $\alpha_1 = 1$  (Fig. 4).

Implications of the above theorems for what concerns the ability of approximating distributions with low coefficient of variation are drawn in [6].

3.3. The minimum coefficient of variation of scaled DPH distributions

For scaled DPH distribution Theorem 3 can be restated as follows.

**Theorem 4.** *The  $cv_{\min}^2$  of a scaled DPH r.v. of order  $n$  with scale factor  $\delta$  and mean  $m_\tau = \delta m_u$  is*

$$\frac{\langle m_\tau / \delta \rangle (1 - \langle m_\tau / \delta \rangle)}{(m_\tau / \delta)^2} \quad \text{if } m_\tau \leq n\delta, \quad \frac{1}{n} - \frac{\delta}{m_\tau} \quad \text{if } m_\tau > n\delta. \tag{5}$$




Fig. 4.  $m_u > n$ .

The scaled DPH r.v. which exhibits the  $cv_{\min}^2$  has the same structure of Figs. 3 and 4, as in the unscaled case (see Theorem 3).

**Corollary 2.** For finite mean  $m_\tau$ , as  $\delta \rightarrow 0$  only the second part of (5) remains effective, and  $cv_{\min}^2 \rightarrow 1/n$  as  $\delta \rightarrow 0$ .

Corollary 2 proves that the  $cv_{\min}^2$  of the DPH class converges to the  $cv_{\min}^2$  of the CPH class of the same order as  $\delta \rightarrow 0$ . The following corollary presents a stronger convergence result for the case of approximating distributions with low coefficient of variation. It is about the convergence of the distributions. In the corollary and its proof the term *best cv fitting* PH approximation will be used which refers to the PH approximation that exhibits the same mean and provides the closest approximation for the second moment.

**Corollary 3.** For distributions whose coefficient of variation is lower than the lowest coefficient of variation attainable by PH distributions, the best cv fitting scaled DPH approximation converges, in distribution, to the best cv fitting CPH approximation of the same distribution as  $\delta$  tends to 0.

**Proof.** Both the CPH and the DPH classes have limits in approximating distributions with low coefficient of variation. As given in Theorems 2 and 3, the best cv fitting approximation of a distribution with coefficient of variation less than these limits is the Erlang( $n$ ) distribution in both the continuous and the discrete case (assuming that  $m_\tau > n\delta$ ).

The representation  $(\alpha, \mathbf{Q})$  of the continuous Erlang( $n$ ) with mean  $m_\tau$  and the representation  $(\alpha, \mathbf{B})$  of the discrete Erlang( $n$ ) with mean  $m_\tau$ , scale factor  $\delta$  are

$$\alpha = \{1, 0, \dots, 0\}, \quad \mathbf{Q} = \begin{bmatrix} -\frac{n}{m_\tau} & \frac{n}{m_\tau} & 0 & \dots & 0 \\ 0 & -\frac{n}{m_\tau} & \frac{n}{m_\tau} & \dots & \\ & & \ddots & & \\ 0 & & & \dots & -\frac{n}{m_\tau} \end{bmatrix},$$

$$\alpha = \{1, 0, \dots, 0\}, \quad \mathbf{B} = \begin{bmatrix} 1 - \frac{n\delta}{m_\tau} & \frac{n\delta}{m_\tau} & 0 & \dots & 0 \\ 0 & 1 - \frac{n\delta}{m_\tau} & \frac{n\delta}{m_\tau} & \dots & \\ & & \ddots & & \\ 0 & & & \dots & 1 - \frac{n\delta}{m_\tau} \end{bmatrix}.$$

Note that  $\mathbf{B} = \mathbf{I} - \mathbf{Q}\delta$  and Corollary 3 follows from Corollary 1. □

In this particular case, when the structure of the best cv fitting scaled DPH and CPH distributions are known, we can show that the distribution of the best cv fitting scaled DPH distribution converges to the




Fig. 5. DPH representation of the discrete uniform distribution  $[a = 2, b = 6]$ .

distribution of the best cv fitting CPH distribution when  $\delta \rightarrow 0$ . Unfortunately, the same convergence property cannot be proved in general, since the structural properties of the best cv fitting PH distributions are not known and they depend on the chosen (arbitrary) optimization criterion. Instead, in Section 4 we provide an extensive experimental study on the behavior of the approximating scaled DPH and CPH distributions as a function of the scale factor  $\delta$ .

### 3.4. DPH distributions with finite support

Another peculiar characteristic of the DPH class is to contain distributions with finite support. A DPH distribution has finite support if its structure does not contain cycles and self-loops (any cycle or self-loop implies an infinite support).

Let  $[a, b]$  be the finite support of a given distribution, with  $a, b \geq 0$  and  $a \leq b$  (when  $a = b$  the finite support distribution reduces to a deterministic distribution with mass 1 at  $a = b$ ). If  $a/\delta$  and  $b/\delta$  are both integers, it is possible to construct a scaled DPH of order  $b/\delta$  for which the probability mass function has non-zero elements only for the values  $a, a + \delta, a + 2\delta, \dots, b$ . As an example, the discrete uniform distribution between  $a = 2$  and  $b = 6$  is reported in Fig. 5, for scale factor  $\delta = 1$ .

## 4. The optimal $\delta$ in PH fitting

The scale factor  $\delta$  provides a new degree of freedom in fitting, and, furthermore, since the limit of a DPH distribution for  $\delta \rightarrow 0$  is a CPH distribution, the optimization of the scale factor in a fitting problem provides a quantitative way to decide whether a continuous or a discrete approximation performs better in the given problem. Hence, assuming  $\delta$  as a decision variable, we can consider the CPH and the DPH class as a unique model set in which the choice among DPH or CPH classes is given by the optimal value of  $\delta$ .

Let  $X$  be the continuous r.v. to be fit by a PH distribution, and let  $F_X(x)$  be its c.d.f.,  $E(X^i)$  the  $i$ th moment and  $\text{cv}^2(X)$  the squared coefficient of variation. In order to define a fitting procedure, a distance measure between  $X$  and the approximating PH distribution needs to be defined. Then, the fitting algorithm provides the PH distribution which minimizes the chosen distance measure. The minimization of the measure is performed as follows. Starting from the initial guess, the non-linear optimization problem is solved by an iterative linearization method. In each step the partial derivatives with respect to the parameters of the PH distribution are computed numerically. Then, the simplex method is applied to determine the direction in which the distance measure decreases optimally. The constraints of the linear programming are given by probabilistic constraints, and by confining the change of parameters. (More detailed description of the fitting procedure, which is implemented in the tool PhFit [16], is given in [15].)

In order to compare, in a unified framework, the goodness of the approximation reached by CPH and DPH distributions, we need to choose a distance measure that is meaningful and applicable both in the continuous as well as in the discrete setting. The selected distance measure is the squared area difference

between the original c.d.f.  $F(\cdot)$  and the approximating c.d.f.  $\hat{F}(\cdot)$ :

$$D = \int_x (F(x) - \hat{F}(x))^2 dx. \tag{6}$$

The distance measure  $D$  is easily applicable for any combination of discrete and continuous distributions. All the numerical experiments reported in the sequel are based on the minimization of the area difference given in (6).

4.1. Fitting distributions with low  $cv^2$

The following considerations provide practical upper and lower bounds to guide in the choice of a suitable scale factor  $\delta$ , and are mainly based on the dependence of the minimal coefficient of variation of a scaled DPH distribution on the order  $n$  and the mean  $m_\tau$ .

Since we only consider DPH distributions with no mass at zero, the mean of any unscaled DPH distribution is greater than 1. This means that  $\delta$  should be less than  $E(X)$ . However, a more convenient upper bound that exploits the flexibility associated with the  $n$  phases is given by

$$\delta \leq \frac{E(X)}{n - 1}. \tag{7}$$

If the squared coefficient of variation of the distribution to be approximated is less than  $1/n$ ,  $\delta$  should satisfy the following relation (see Theorem 3):

$$\delta > \left( \frac{1}{n} - cv^2(X) \right) E(X). \tag{8}$$

Let  $X$  be a lognormal r.v. with parameters (1, 0.2), whose mean is  $E(X) = 1$  and  $cv^2(X) = 0.0408$  (this distribution is the distribution L3 taken from the benchmark examined in [6,9], hence we refer to it as L3). Table 1 reports the lower and upper bounds of  $\delta$ , with  $n = 2, 4, 8, 12$ , computed from (8) and (7).

The c.d.f. and p.d.f. of the approximating CPH and DPH distributions of order  $n = 10$ , with different scale factors  $\delta$ , are presented in Fig. 6. When considering the approximate DPH distribution, the  $f(x)$  values are calculated at the discrete points  $(\delta, 2\delta, 3\delta, \dots, k\delta, \dots)$  to which the following mass is assigned:

$$f(k\delta) = \frac{1}{\delta} (F(k\delta) - F((k - 1)\delta)). \tag{9}$$

For the ease of visual interpretation the points are connected with a line.

Table 1  
Upper and lower bound of  $\delta$  for fitting distribution L3

$n$	Lower bound of $\delta$ (Eq. (8))	Upper bound of $\delta$ (Eq. (7))
4	0.2092	0.333
8	0.0842	0.1428
12	0.0425	0.0909
16	0.0217	0.0666




Fig. 6. Approximating the L3 distribution with 10-phase PH approximations.

When  $\delta$  is less than its lower bound the required  $cv^2$  cannot be attained; when  $\delta$  becomes too large the wide separation of the discrete steps increases the approximation error; when  $\delta$  is in the proper range (e.g.  $n = 10$ ,  $\delta = 0.06$ ) a reasonably good fit is achieved. This example also suggests that an optimal value of  $\delta$  exists that minimizes the chosen distance measure  $D$  in (6).

In order to display the goodness of fit for the L3 distribution, Fig. 7 shows the distance measure  $D$  as a function of  $\delta$  for various values of the order  $n$ . A minimum value of  $D$  is attained in the range where the parameters fit the bounds of Table 1. Notice also that, as  $\delta$  increases, the advantage of having more phases disappears, according to Theorem 4. The circles in the left part of this figure (as well as in all the successive figures) indicate the corresponding distance measure  $D$  obtained from CPH fitting. The figure (and the subsequent ones as well) suggests that the distance measure obtained from DPH fitting converges to the distance measure obtained by the CPH approximation as  $\delta$  tends to 0.

The lowest applied scale factor is 0.01. Lower values of the scale factor renders the fitting procedure numerically unstable because the diagonal elements of the transition matrix are close to 1.




Fig. 7. Distance measure as the function of the scale factor  $\delta$  for low  $cv^2$  (L3).

#### 4.2. Fitting distributions with high $cv^2$

We have seen in the previous subsections that it is beneficial to approximate distributions with a low coefficient of variation by means of a DPH distributions. In this section, we investigate the optimal value of  $\delta$  when fitting distributions with a high coefficient of variation.

Let  $X$  be a lognormal r.v. with parameters  $(1, 1.8)$  (this is the distribution L1 taken from the benchmark in [6,9]). For  $X$  we have  $E(X) = 1$  and  $cv^2(X) = 24.534$ . Fig. 8 shows the measure of the goodness of fit as a function of  $\delta$  for various orders  $n$  (the cases when the number of phases is greater than 2 result in the same goodness of fit). The distance measures  $D$  decreases as  $\delta \rightarrow 0$  indicating that the optimal fitting is achieved by applying CPH distribution. This example suggests that, for distributions with infinite support and  $cv^2(X) > 1/n$ , the optimal value of  $\delta$  tends to 0, implying that the best-fit is obtained by a CPH. However, this conclusion might not be true for distributions with finite support, this case is explored in the next section.




Fig. 8. Distance measure as the function of the scale factor  $\delta$  for high  $cv^2$  (L1).




Fig. 9. Distance measure as the function of the scale factor  $\delta$  for Uniform(1, 2) (U2).

4.3. Fitting distributions with finite support

Two features must be considered, namely the  $cv^2$  and the maximum value of the finite support. It should be stressed that the chosen distance measure  $D$  in (6) can be considered as not completely appropriate in the case of finite support, since it does not force the approximating PH to have its mass confined in the finite support and 0 outside.

Let  $X$  be a uniform r.v. over the interval  $[1, 2]$ , with  $E(X) = 1.5$  and  $cv^2(X) = 0.0370$  (this is the distribution U2 taken from the benchmark in [6,9]). Fig. 9 shows the distance measure as a function of  $\delta$  for various orders  $n$ . It is evident that, for each  $n$ , a minimal value of  $\delta$  is obtained, that provides the best approximation according to the chosen distance measure.

As a second example, let  $X$  be a uniform r.v. over the interval  $[0, 1]$ , with  $E(X) = 0.5$  and  $cv^2(X) = 0.333$  (this is the distribution U1 taken from the benchmark in [6,9]). Fig. 10 shows the distance measure




Fig. 10. Distance measure as the function of the scale factor  $\delta$  for Uniform(0, 1) (U1).

as a function of  $\delta$  for various orders  $n$ . Since, in this example  $cv^2(X) = 0.333$ , an order  $n = 3$  is large enough for a CPH to attain the coefficient of variation of the distribution. Nevertheless, the optimal  $\delta$  in Fig. 10, which minimizes the distance measure  $D$  for high order PH ( $n > 2$ ), ranges between  $\delta = 0.02$  and  $0.05$ , thus leading to the conclusion that a DPH provides a better fit. This example evidences that the coefficient of variation is not the only factor which influences the optimal  $\delta$  value. The shape of the distribution plays an essential role as well. Our experiments show that a discontinuity in the p.d.f. (or in the c.d.f.) is hard to approximate with CPH, hence in the majority of these cases DPH provides a better approximation.

Fig. 11 shows the c.d.f. and the p.d.f. of the U1 distribution, compared with the best-fit PH approximations of order  $n = 10$ , and various scale factors  $\delta$ . In the case of DPH approximation, the  $f(x)$  values are calculated as in (9). With respect to the chosen distance measure, the best approximation is obtained for  $\delta = 0.03$ , which corresponds to a DPH distribution with infinite support. When  $\delta = 0.1$  the approximate distribution has a finite support. Hence, the value  $\delta = 0.1$  (for  $n = 10$ ) provides a DPH able to represent the property that the r.v. is less than 1. Another fitting criterion may, of course, stress this property.




Fig. 11. Approximating the Uniform(0, 1) distribution (U1).




Fig. 12. Distance measure as the function of the scale factor for the Exponential(1.0) distribution.

#### 4.4. Approximating exponential distributions

When it comes to discrete approximation of continuous models, exponential durations are usually approximated by a discrete PH distribution with a single phase (see, for example, [14]). This approximation, however, is precise only if the scale factor is small enough. Fig. 12 shows the goodness of fitting as function of the scale factor for the exponential distribution with mean equal to 1.0. The figure suggests that it is not worth to use more than two phases to approximate exponential distributions. (Note that this statement might not be true in case of other distance measures.)

### 5. Approximating non-Markovian models

Section 4 has explored the problem of how to find the best-fit among either a DPH or a CPH distribution by tuning the scale factor  $\delta$ . When dealing with a stochastic model of a system that incorporates non-exponential distributions, a well-known solution technique consists in a Markovianization of the underlying non-Markovian process by substituting the non-exponential distribution with a best-fit PH distribution, and then expanding the state space. A natural question arises also in this case, on how to decide between discrete (using DPH) and continuous (using CPH) approximation, in order to minimize the error in the performance measures we are interested in for the overall model.

One possible way to handle this problem could consist in finding the best PH fit for any single distribution and to plug them in the model. In the present paper, we only consider the case where the PH distributions are either all discrete with the same scale factor or they are all continuous. This is clearly a limitation in choosing the scale factor because a single scale factor is chosen for the whole model. Various embedding techniques have been explored in the literature for mixing DPH with different scale factors and CPH [13,19], but these techniques are out of the scope of the paper.

In order to quantitatively evaluate the influence of the scale factor on some performance measures defined at the system level, we have considered a preemptive M/G/1/2/2 queue with two classes of customers. We have chosen this example because accurate analytical solutions are available both in




Fig. 13. The state space of the considered M/G/1/2/2 queue.

transient condition and in steady state using the methods presented in [13]. The general distribution  $G$  is taken from the set of distributions (L1, L3, U1, U2) already considered in the previous section.

Customers arrive at the queue with rate  $\lambda = 0.5$  in both classes. The service time of a higher priority job is exponentially distributed with parameter  $\mu = 1$ . The service time distribution of the lower priority job is either L1, L3, U1 or U2. Arrival of a higher priority job preempts the lower priority one. The policy associated to the preemption of the lower priority job is preemptive repeat different (prd), i.e. after the departure of the higher priority customer the service of the low priority customer starts from the beginning with a new service time sample.

The system has four states (Fig. 13): in state  $s_1$  the server is empty, in state  $s_2$  a higher priority customer is under service with no lower priority customer in the system, in state  $s_3$  a higher priority customer is under service with a lower priority customer waiting, in state  $s_4$  a lower priority job is under service (in this case there cannot be a higher priority job).

### 5.1. Steady-state behavior

Let  $p_i$  ( $i = 1, \dots, 4$ ) denote the steady-state probability of the M/G/1/2/2 queue obtained from an exact analytical solution.




Fig. 14.  $\epsilon_{SUM}$  for distribution L3 with 1-phase approximation for exponential durations.




Fig. 15.  $\epsilon_{SUM}$  for distribution L3 with 2-phase approximation for exponential durations.

In order to evaluate the correctness of the PH approximation we have solved the model by substituting the original general distribution (either L1, L3, U1 or U2) with approximating DPH or CPH distributions. (The fitting was performed applying the same distance measure as before, it is given in (6)). Let  $\hat{p}_i$  ( $i = 1, \dots, 4$ ) denote the steady-state probability of the M/PH/1/2/2 queue with the PH approximation.

The overall approximation error is measured in terms of the difference between the exact steady-state probabilities  $p_i$  and the approximate steady-state probabilities  $\hat{p}_i$ . Two error measures are defined as:

$$\epsilon_{SUM} = \sum_i \frac{|p_i - \hat{p}_i|}{p_i} \quad \text{and} \quad \epsilon_{MAX} = \max_i \frac{|p_i - \hat{p}_i|}{p_i}.$$

All the experiments were performed applying both 1-phase and 2-phase approximation for the exponential durations. The evaluated numerical values for  $\epsilon_{SUM}$  and  $\epsilon_{MAX}$  are reported in Figs. 14 and 15 and Figs. 16 and 17 for the distribution L3. Since the behavior of  $\epsilon_{MAX}$  is very similar to the behavior of  $\epsilon_{SUM}$  in all




Fig. 16.  $\epsilon_{MAX}$  for distribution L3 with 1-phase approximation for exponential durations.




Fig. 17.  $\epsilon_{MAX}$  for distribution L3 with 2-phase approximation for exponential durations.

the cases, for the other cases we report  $\epsilon_{SUM}$  only (L1: Figs. 18 and 19, U1: Figs. 20 and 21, U2: Figs. 22 and 23).

Figs. 24 and 25 report two cases when both the service times are non-exponential. In the first case both the service distributions are L3, while in the second case, higher (lower) priority customers are served according to distribution L3 (L1).

Based on the above experiments, it is rather hard to draw general guidelines for the choice of the scale factor. Some observations, however, can be made:

- If all the optimal scale factors are 0, the approximation error in the evaluation of the performance indices of the global model can be minimized by resorting to continuous PH type approximation. See, for example, the case when the only non-exponential distribution in the model is L1 (Fig. 18).
- When some of the individual optimal scale factors are 0, while others are not (this is the case, for example, when there are exponential durations and durations with low coefficient of variation in the




Fig. 18.  $\epsilon_{SUM}$  for distribution L1 with 1-phase approximation for exponential durations.




Fig. 19.  $\epsilon_{SUM}$  for distribution L1 with 2-phase approximation for exponential durations.

same model, Fig. 14), there is an optimal scale factor value (or a range) that gives optimal results for what concerns performance measures of the global model. In this case, on the one hand, decreasing the scale factor worsen the precision of the approximation because with lower scale factor durations with low coefficient of variation are approximated worse. On the other hand, increasing the scale factor worsen the approximation because the approximation of the exponential durations are getting less accurate (see Fig. 14).

- In the cases when one of the individual optimal scale factors is rather far from 0, there is a wide range of scale factor values for which the discrete approximation yields better results than the continuous one. See, for example, Fig. 22 where in case of 10 phases the discrete approximation gives better results in the range [0, 0.2].
- Applying 2-phase discrete PH approximations for the exponential durations slows down the rate at which the approximation error grows with increasing scale factor. See, for example, Fig. 14 compared to Fig. 15.




Fig. 20.  $\epsilon_{SUM}$  for distribution U1 with 1-phase approximation for exponential durations.




Fig. 21.  $\epsilon_{\text{SUM}}$  for distribution U1 with 2-phase approximation for exponential durations.

## 5.2. Transient behavior

In order to investigate the approximation error in the transient behavior, we have considered distribution U2 for the service time and we have computed the transient probability of state  $s_1$  with two different initial conditions. Fig. 26 depicts the transient probability of state  $s_1$  when the process is started in state  $s_1$  with probability 1. Fig. 27 depicts the transient probability of the same state,  $s_1$ , when the service of a lower priority job starts at time 0 (the process is started with probability 1 in  $s_4$ ). All approximations are with DPH distributions of order  $n = 10$ . Only the DPH approximations are depicted because the CPH approximation is very similar to the DPH one with scale factor  $\delta = 0.03$ . In the first case (Fig. 26), the scale factor  $\delta = 0.03$ , which was the optimal one from the point of view of fitting the single distribution in isolation, provides the most accurate results for the transient analysis as well. Instead, in the second case, the approximation with a scale factor  $\delta = 0.2$  captures better the sharp change in the transient




Fig. 22.  $\epsilon_{\text{SUM}}$  for distribution U2 with 1-phase approximation for exponential durations.




Fig. 23.  $\epsilon_{SUM}$  for distribution U2 with 2-phase approximation for exponential durations.




Fig. 24.  $\epsilon_{SUM}$  for service times L3–L3 with 1-phase approximation for exponential durations.






Fig. 25.  $\epsilon_{SUM}$  for service times L1–L3 with 1-phase approximation for exponential durations.

Fig. 26. Transient probabilities starting from  $s_1$ .Fig. 27. Transient probabilities starting from  $s_4$ .

probability. Moreover, this value of  $\delta$  is the only one among the values reported in the figure that results in zero probability for time points smaller than 1. In other words, the second example depicts the advantage given by DPH distributions to model durations with finite support. This example suggests also that DPH approximation can be of importance when preserving reachability properties is crucial (like in modeling time-critical systems) and, hence, DPH approximation can be seen as a bridge between the world of stochastic modeling and the world of functional analysis and model checking [5].

## 6. Concluding remarks

The main result of this paper is that, with the notion of the scale factor, we gain a unified view of the DPH and the CPH distributions. The optimal value of the scale factor  $\delta$ ,  $\delta_{opt}$ , determines the best

distribution in a fitting experiment. When  $\delta_{\text{opt}} = 0$  the best choice is a CPH distribution, while when  $\delta_{\text{opt}} > 0$  the best choice is a DPH distribution. This paper has also shown that the transition from DPH class to CPH class is continuous with respect to several properties, like the distance (denoted by  $D$  in (6)) between the original and the approximate distributions. The paper presents limit theorems for special cases; however, extensive numerical experiments show that the limiting behavior is far more general than the special cases considered in the theorems.

The numerical examples have also evidenced that for very small values of  $\delta$ , the diagonal elements of the transition probability matrix are very close to 1, rendering numerically unstable the DPH fitting procedure.

A deep analytical and numerical sensitivity analysis is required to draw more general conclusions for the “optimal  $\delta$  value” at the mode level. It is definitely a field of further research.

Finally, we summarize the advantages and the disadvantages of applying approximate DPH models (even with optimal  $\delta$  value) with respect to using CPH approximations.

### 6.1. Advantages of using DPH

An obvious advantage of the application of DPH distributions is that one can have a closer approximation of distributions with low coefficient of variation. Another important quantitative property of the DPH class is that it can capture distributions with finite support and deterministic values. This property allows to capture the periodic behavior of a complex stochastic model, while any CPH-based approximation of the same model tends to a steady state.

Numerical experiments have also shown that DPH can better approximate the distributions with some abrupt or sharp changes in the c.d.f. or in the p.d.f.

### 6.2. Disadvantages of using DPH

There is a definite disadvantage of discrete time approximation of continuous time models. In the case of CPH approximation, coincident events do not have to be considered (they have zero probability of occurrence). Instead, when applying DPH approximation coincident events have to be handled, and their consideration may burden significantly the complexity of the analysis.

## Acknowledgements

This work has been performed under the Italian-Hungarian R&D program supported by the Italian Ministry of Foreign Affairs and the Hungarian Ministry of Education. A. Bobbio and A. Horváth were partially supported by the MIUR under the project FIRB-Perf; M. Telek was partially supported by Hungarian Scientific Research Fund (OTKA) under Grant No. T-34972.

## References

- [1] D. Aldous, L. Shepp, The least variable phase-type distribution is Erlang, *Stoch. Mod.* 3 (1987) 467–473.
- [2] S. Asmussen, O. Nerman, Fitting phase-type distributions via the EM algorithm, in: *Proceedings of the Symposium i Advent Statistik*, Copenhagen, 1991, pp. 335–346.

- [3] A. Bobbio, A. Cumani, ML estimation of the parameters of a PH distribution in triangular canonical form, in: G. Balbo, G. Serazzi (Eds.), *Computer Performance Evaluation*, Elsevier, Amsterdam, 1992, pp. 33–46.
- [4] A. Bobbio, A. Horváth, Model checking time Petri nets using NuSMV, in: *Proceedings of the Fifth International Workshop on Performability Modeling of Computer and Communication Systems (PMCCS5)*, 2001, pp. 100–104.
- [5] A. Bobbio, A. Horváth, Petri nets with discrete phase-type timing: a bridge between stochastic and functional analysis, *Electronic Notes in Theoretical Computer Science* 52 (3) (2001). <http://www.elsevier.nl/locate/entcs/volume52.html>.
- [6] A. Bobbio, A. Horváth, M. Scarpa, M. Telek, Acyclic discrete phase type distributions: properties and a parameter estimation algorithm, *Perform. Eval.* 54 (1) (2003) 1–32.
- [7] A. Bobbio, A. Puliafito, M. Telek, A modeling framework to implement preemption policies in non-Markovian SPN, *IEEE Trans. Softw. Eng.* 26 (2000) 36–54.
- [8] A. Bobbio, A. Puliafito, M. Telek, K. Trivedi, Recent developments in non-Markovian stochastic Petri nets, *J. Syst. Circuits Comput.* 8 (1) (1998) 119–158.
- [9] A. Bobbio, M. Telek, A benchmark for PH estimation algorithms: results for acyclic-PH, *Stoch. Mod.* 10 (1994) 661–677.
- [10] G. Ciardo, R. German, C. Lindemann, A characterization of the stochastic process underlying a stochastic Petri net, *IEEE Trans. Softw. Eng.* 20 (1994) 506–515.
- [11] A. Cumani, On the canonical representation of homogeneous Markov processes modelling failure-time distributions, *Microelectr. Reliab.* 22 (1982) 583–602.
- [12] A. Cumani, Esp—A package for the evaluation of stochastic Petri nets with phase-type distributed transition times, in: *Proceedings of the International Workshop on Timed Petri Nets*, Torino, Italy, 1985, pp. 144–151.
- [13] R. German, *Performance Analysis of Communication Systems: Modeling with non-Markovian Stochastic Petri Nets*, Wiley, New York, 2000.
- [14] A. Horváth, A. Puliafito, M. Scarpa, M. Telek, Analysis and evaluation of non-Markovian stochastic Petri nets, in: *Proceedings of the 11th Performance TOOLS*, Schaumburg, IL, USA, March 2000, *Lecture Notes in Computer Science*, vol. 1786, Springer, Berlin, pp. 171–187.
- [15] A. Horváth, M. Telek, Approximating heavy tailed behavior with phase type distributions, in: *Proceedings of the Third International Conference on Matrix-Analytic Methods in Stochastic Models*, Leuven, Belgium, June 2000.
- [16] A. Horváth, M. Telek, Phfit: A general phase-type fitting tool, in: *Proceedings of the 12th Performance TOOLS*, Imperial College, London, April 2002, *Lecture Notes in Computer Science*, vol. 2324, Springer, Berlin.
- [17] A. Horváth, M. Telek, Time domain analysis of non-Markovian stochastic petri nets with pri-transitions, *IEEE Trans. Softw. Eng.* 28 (20) (2002) 933–943.
- [18] M. Johnson, M. Taaffe, Matching moments to phase distributions: nonlinear programming approaches, *Stoch. Mod.* 6 (1990) 259–281.
- [19] R. Jones, G. Ciardo, On phased delay stochastic Petri nets: Definition and application, in: *Proceedings of the Ninth International Workshop on Petri Nets and Performance Models (PNPM01)*, IEEE Computer Society, 2001.
- [20] A. Lang, J.L. Arthur, Parameter approximation for phase-type distributions, in: *Matrix-analytic Methods in Stochastic Models*, *Lecture Notes in Pure and Applied Mathematics*, Marcel Dekker, New York, 1996, pp. 151–206.
- [21] C. Lindemann, DSPNexpress: a software package for the efficient solution of deterministic and stochastic Petri nets, *Perform. Eval.* 22 (1995) 3–21.
- [22] M. Neuts, Probability distributions of phase type, in: *Liber Amicorum Prof. Emeritus H. Florin*, University of Louvain, 1975, pp. 173–206.
- [23] M. Neuts, *Matrix Geometric Solutions in Stochastic Models*, Johns Hopkins University Press, Baltimore, MD, 1981.
- [24] L. Schmickler, MEDA: mixed Erlang distributions as phase-type representations of empirical distribution functions, *Stoch. Mod.* 8 (1992) 131–156.
- [25] M. Telek, Minimal coefficient of variation of discrete phase type distributions, in: *Proceedings of the Third International Conference on Matrix-Analytic Methods in Stochastic Models (MAM3)*, Leuven, Belgium, 2000, pp. 391–400.



**A. Bobbio** graduated in nuclear engineering from Politecnico di Torino in 1969. In 1971 he joined the Istituto Elettrotecnico Nazionale Galileo Ferraris di Torino, where he was involved in activities connected with the organization of a “Reliability Club” (Circolo dell’Affidabilità). In 1992, he became associate professor at the Department of “Elettronica per l’Automazione” of the University of Brescia, and in 1995 he moved to the Department of Informatica of the University of Torino. In 2000, he became full professor at the Università del Piemonte Orientale of Alessandria, Italy. His activity is mainly focused on the modeling and analysis of the performance and reliability of stochastic systems, with particular emphasis on Markov chains and stochastic Petri Nets. Bobbio has spent various research periods at the Department of Computer Science of the Duke University (Durham, NC, USA), at the Technical University of Budapest and at the Department of Computer Science and Engineering at the Indian Institute of Technology in Kanpur (India). He has been principal investigator and leader of research groups in various research projects with public and private institutions. He is author of several papers in international journals as well as communications to international conferences and workshops.



**A. Horváth** was born in 1974 in Budapest where he received the M.Sc. degree in computer science from the University of Technology and Economics. From 1998 to 2002 he was a Ph.D. student supervised by Miklós Telek at the same university. From 2003 he is a researcher at the University of Turin (Italy). His research interests are in the area of stochastic processes including performance analysis of non-Markovian systems and modeling issues of communication networks.



**M. Telek** received the M.Sc. degree in electrical engineering from the Technical University of Budapest in 1987. After graduation he joined the Hungarian Post Research Institute where he studied the modelling, analysis and planning aspects of communication networks. Since 1990 he has been with the Department of Telecommunications of the Technical University of Budapest, where he is an associate professor now. He received the candidate of science degree from the Hungarian Academy of Science in 1995. His current research interest includes stochastic performance modeling and analysis of computer and communication systems.